

ULTRASONIC- AND MICROWAVE-ASSISTED EXTRACTION OF
CURCUMINOIDS AND CYCLODEXTRIN COMPLEXES OF CURCUMIN
FROM *C. domestica* Val.

BINTA HADI JUME

A thesis submitted in fulfilment of the
requirements for the award of the degree of
Doctor of Philosophy(Chemistry)

Faculty of Science
Universiti Teknologi Malaysia

MAY 2014

I dedicated this thesis to my parent and my family for their support, encouragement and understanding.

ACKNOWLEDGEMENTS

I wish to express my deep and sincere gratitude to Professor Dr. Mohd Marsin Sanagi, my main supervisor and mentor during PhD study, for his helpful support, advice, constructive criticism, and critical review of the thesis. Special appreciation is extended to my Co-supervisors Professor Dr. Wan Aini Wan Ibrahim and Dr. Shajarahtunnur Jamil for their help, support, and helpful technical discussions. I would also like to thank Professor Dr. Madzlan Bin Aziz and Professor Dr. Wan Azelee Wan Abu Bakar, Dean of Faculty of Science and Head of Department of Chemistry, Universiti Teknologi, Malaysia, respectively, for their guidance and feedback. I also appreciate Professor R. M. Smith and Professor Hassan Aboul-Enein for their valuable help and guidance.

I would like to express my deep appreciation to my beloved father Alhaji Hadi Jume. He sacrificed a lot to raise me and to support my education. I thank my devoted mother Hajiya Sa'adatu Hadi Jume from the bottom of my heart for her prayers and encouragement. I will never forget my beloved husband Dr. Mu'azu Mohammed Abdullahi, for his help, support, prayers and encouragement. He was always there to help and support me at all the times. He sacrificed a lot to provide me comfort and a good study environment during this PhD study. My special appreciation is also extended to my little son Al-Amin Mohammed Mu'azu and my baby girls; Faridah Mohammed Mu'azu and Fadilah Mohammed Mu'azu. I forgot my troubles and found peace of mind when I'm playing with them. Foremost, I also give thanks to Almighty Allah for giving me the opportunity to meet and study with these great people at UTM. I cherish my entire friends for their support in one way or the other.

ABSTRACT

This study investigates the isolation of curcuminoids from *Curcuma domestica* Val. using ultrasonic-assisted extraction (UAE), microwave-assisted extraction (MAE) compared with conventional cold solvent extraction method and the use of inclusion complexation of curcumin with methyl- β -cyclodextrin (M β -CD) for improving their solubility. The extractions were optimized by determining the content of three curcuminoid markers, namely curcumin (C), demethoxycurcumin (DMC) and bisdemethoxycurcumin (BDMC). The extraction efficiencies were compared in terms of extraction time, sample through-put and solvent consumption. The optimized parameters for UAE of curcuminoids were extraction amplitude of 100, particle size of 0.30-0.60 mm, extraction time of 20 min, extraction solvent volume of 10 mL and extraction temperature of 60°C. Meanwhile, the relative recoveries (RRs) for aqueous extraction were in the range of 91.59-98.99%, 89.79-94.95% and 89.33-94.77% for C, DMC and BDMC, respectively. Although UAE using methanol resulted in a slightly higher extraction yield and shorter extraction time compared to those using water, both methods showed similar pattern of results. Despite these, water is cost effective, safe and environmentally friendly. These advantages of aqueous solvent can be used as a yard stick to substitute organic solvents for UAE of curcuminoids from *C. domestica*. At maximum set energy, the MAE optimum extraction parameters were particle size of 0.30-0.60 mm, extraction time of 3 min, extraction solvent volume of 10 mL and extraction temperature of 60°C with RRs of 92.48-99.44%, 90.58-97.43% and 90.03-96.07% for C, DMC and BDMC, respectively. Both UAE and MAE applications showed remarkable improvements in terms of extraction time, solvent consumption, extraction yield and the quality of extracts compared to conventional cold solvent extractions method. However, as compared to UAE, the optimized MAE application was better in term of quantity of curcuminoids. MAE is also simpler, faster, more efficient approach and allows the possibility of simultaneous multiple extractions. The inclusion complex formed using M β -CD with the application of MAE was more stable than that with UAE based on the stability constant (K_C) values of 213.08 M⁻¹ and 515.19 M⁻¹, for UAE and MAE, respectively. Results from characterization of the inclusion complex with scanning electron microscope showed that co-precipitation method was best for UAE while all of the mixing methods can be used for the inclusion complexation with MAE application. The kneading and co-precipitation methods were found to be the best for the inclusion complexation between turmeric rhizome oleoresins and M β -CD in UAE, while all of the mixing methods were found to be suitable for inclusion complexation of turmeric rhizome oleoresins with M β -CD in MAE as indicated by Fourier transform infrared spectroscopy.

ABSTRAK

Kajian ini menyelidik pengasingan kurkumin daripada *Curcuma domestica* Val. menggunakan pengekstrakan berbantuan ultrasonik (UAE) dan pengekstrakan berbantuan gelombang mikro (MAE) berbanding dengan kaedah pengekstrakan pelarut sejuk konvensional dan penggunaan pengkompleksan rangkuman kurkumin dengan metil- β -siklodekstrin (M β -CD) bagi meningkatkan keterlarutan. Pengekstrakan telah dioptimumkan dengan menentukan kandungan tiga kurkuminoid penanda iaitu kurkumin (C), demetoksikurkumin (DMC) dan bisdemetoksikurkumin (BDMC). Kecekapan pengekstrakan telah dibandingkan daripada segi masa pengekstrakan, kadar pemindahan sampel dan penggunaan pelarut. Parameter optimum bagi UAE kurkuminoid ialah amplitud pengekstrakan 100, saiz partikel 0.30-0.60 mm, masa pengekstrakan 20 min, isipadu pelarut pengekstrak 10 mL dan suhu pengekstrakan 60°C. Sementara itu, pengembalian relatif (RR) bagi pengekstrakan akueus adalah dalam julat 91.59-98.99%, 89.79-94.95% and 89.33-94.77% bagi masing-masing C, DMC dan BDMC. Walaupun UAE yang menggunakan metanol menghasilkan pengembalian pengekstrakan yang lebih tinggi sedikit dan masa pengekstrakan lebih singkat berbanding dengan pengekstrakan yang menggunakan air, kedua-dua kaedah ini menunjukkan pola hasil yang sama. Di sebalik itu, air adalah efektif kos, selamat dan mesra alam. Kelebihan pelarut akueus ini boleh digunakan sebagai kayu pengukur untuk menggantikan pelarut organik bagi UAE kurkuminoid daripada *C. domestica*. Pada tenaga maksimum yang ditetapkan, parameter optimum MAE ialah saiz partikel 0.30-0.60 mm, masa pengekstrakan 3 min, isipadu pelarut pengekstrakan 10 mL dan suhu pengekstrakan 60°C dengan nilai RR 92.48-99.44%, 90.58-97.43% dan 90.03-96.07% masing-masing bagi C, DMC dan BDMC. Kedua-dua aplikasi UAE dan MAE menunjukkan penambahbaikan yang ketara daripada segi masa pengekstrakan, penggunaan pelarut, hasil pengekstrakan dan kualiti ekstrak berbanding dengan kaedah pengekstrakan pelarut sejuk konvensional. Walau bagaimanapun, berbanding UAE, aplikasi MAE yang optimum adalah lebih baik daripada segi kuantiti kurkuminoid. MAE juga merupakan pendekatan yang lebih ringkas, cepat, lebih cekap dan memungkinan pengekstrakan berganda serentak. Kompleks rangkuman yang dihasilkan menggunakan M β -CD dengan MAE adalah lebih stabil berbanding dengan UAE berdasarkan nilai pemalar kestabilan (K_C) iaitu 213.08 M⁻¹ and 515.19 M⁻¹, bagi masing-masing UAE dan MAE. Hasil daripada pencirian kompleks rangkuman dengan mikroskopi elektron pengimbas menunjukkan kaedah ko-pemendakan adalah terbaik bagi UAE manakala semua ko-pemendakan didapati boleh digunakan bagi pengkompleksan rangkuman dengan penggunaan MAE. Kaedah ulian dan ko-pemendakan didapati terbaik bagi pengkompleksan rangkuman antara oleoresin rizom kunyit dengan M β -CD dalam UAE, manakala kaedah pencampuran yang lain didapati sesuai bagi pengkompleksan rangkuman oleoresin rizom kunyit dengan M β -CD dalam MAE seperti ditunjukkan oleh spektroskopi inframerah transformasi Fourier.

TABLE OF CONTENT

CHAPTER	TITLE	PAGE
	DECLARATION	ii
	DEDICATION	iii
	ACKNOWLEDGEMENTS	iv
	ABSTRACT	v
	ABSTRAK	vi
	TABLE OF CONTENTS	vii
	LIST OF TABLES	xiv
	LIST OF FIGURES	xv
	LIST OF ABBREVIATIONS	xxiv
	LIST OF SYMBOLS	xxvi
	LIST OF APPENDICES	xxvii
1	INTRODUCTION	1
	1.1 Research Background	1
	1.2 Problem Statement	6
	1.3 Objectives of Study	6
	1.4 Scope of the Research	7
	1.5 Justification of the Research	7
	1.6 Framework of the Research	8
	1.7 Thesis Organization	9
2	LITERATURE REVIEW	11
	2.1 Introduction	11

2.2	Usages of Curcumin	12
2.3	Constituents and Chemistry of Turmeric	12
2.4	Extraction of Curcumin	14
2.5	Conventional Extraction Methods	15
2.5.1	Soxhlet Extraction	15
2.5.2	Solvent Extraction	17
2.5.3	Reflux Extraction	17
2.6	Unconventional Extraction Methods	18
2.6.1	Ultrasonic-Assistance Extraction	18
2.6.2	Microwave Extraction	21
2.6.2.1	Mechanism of Microwave Extraction	22
2.6.2.2	Mechanism of Microwave Heating	25
2.6.2.3	Parameters Affecting Microwave-Assisted Extraction	26
2.6.3	Subcritical Water Extraction	28
2.6.4	Supercritical Fluid Extraction System	30
2.6.5	Accelerated Solvent Extraction or Pressurized Liquid Extraction	32
2.7	Structural Features and Characteristics of Cyclodextrin	33
2.7.1	Physico-Chemical Properties of β -Cyclodextrin	35
2.7.2	Applications and Current use of Cyclodextrins	37
2.7.3	Mechanism of Complexation between Cyclodextrin and Insoluble Compound	38
2.7.4	Stoichiometry and Association Constant of Compound-Cyclodextrin Inclusion Complex	39
2.7.5	Method for Determination of Stoichiometry and Association Constant of the Inclusion Complex: The Phase Solubility Method	40
3	EXPERIMENTAL	44
3.1	Introduction	44

3.2	Chemicals and Materials	45
3.3	Extraction of the Components of the Rhizomes of <i>C. domestica</i> Val.	45
3.4	Ultrasonic -Assisted Extraction Procedure	46
3.5	Microwave -Assisted Extraction Procedure	48
3.6	Preparation Stock Solutions	49
3.7	Optimization of High Performance Liquid Chromatography Parameters	49
3.7.1	UV Wavelength Detection	50
3.7.2	Mobile Phase Flow Rate	50
3.7.3	Mobile Phase	50
3.7.4	Temperature of the Column	51
3.7.5	High Performance Liquid Chromatographic Conditions	51
3.8	Cold Solvent Extraction	52
3.9	Optimization of Extraction Parameters and Practical Considerations	52
3.10	Standard Calibration Curve	53
3.11	Method Validation	53
3.11.1	Linearity	54
3.11.2	Limit of Detection	54
3.11.3	Limit of Quantification	55
3.11.4	Relative Recovery	55
3.11.5	Precision	56
3.12.	Phase Solubility	57
3.13	Preparation of Inclusion Complexation in Solution State	58
3.13.1	Kneading Method	59
3.13.2	Co-precipitation Method	59
3.13.3	Physical Mixture	60
3.14	Characterisation of Inclusion Complexes	60
3.14.1	Fourier Transform Infrared Spectroscopy	60
3.14.2	Scanning Electron Microscopy	61

4	ULTRASONIC-ASSISTED EXTRACTION AND COLD SOLVENT EXTRACTION OF CURCUMINOIDS FROM <i>C. domestica</i> Val. USING METHANOL	63
4.1	Introduction	63
4.2	Effect of Extraction Parameters in Ultrasonic-Assisted Extraction of <i>C. domestica</i> Val. with Methanol	64
4.2.1	Effect of Ultrasound Amplitude	65
4.2.2	Effect of Particle Size	66
4.2.3	Effect of Extraction Time	68
4.2.4	Effect of Solvent Volume	70
4.2.5	Effect of Temperature	71
4.2.6	Effect of Methanol Modified with Water	73
4.3	Effect of Optimum Extraction Parameter on Cold Solvent Extraction with Methanol	74
4.3.1	Effect of Extraction Time at Room Temperature	74
4.3.2	Effect of Solvent Volume on the Extraction at Room Temperature	76
4.3.3	Effect of Particle Size on the Extraction at Room Temperature	77
4.4	Validation Method	78
4.4.1	Calibration Curve	79
4.4.2	Limit of Detection and Limit of Quantification	79
4.4.3	Relative Recoveries	79
	4.4.3.1 Methanolic Solvent Extraction with Ultrasound-Assisted Extraction	80
	4.4.3.2 Cold Solvent Extraction with Methanol	80
4.5	High Performance Liquid Chromatograms	81
4.6	Summary	86

5	CYCLODEXTRIN COMPLEXATION ULTRASONIC- ASSISTED EXTRACTION OF CURCUMINOIDS FROM <i>C. domestica</i> Val. USING AQUEOUS SOLVENT	88
5.1	Introduction	88
5.2	Effect of Extraction Parameters on Ultrasonic-Assisted Extraction with Water	89
5.2.1	Effect of Ultrasound Amplitude of Ultrasonic- Assisted Extraction with Water	89
5.2.2	Effect of Particle Size of Sample on the Ultrasonic- Assisted Extraction with Water	91
5.2.3	Effect of Extraction Time on Ultrasonic-Assistance Extraction with Water	93
5.2.4	Effect of Solvent Volume on Ultrasonic-Assistance Extraction with Water	94
5.2.5	Effect of Temperature on the Ultrasonic-Assistance Extraction with Water	96
5.3	Effect of Optimum Extraction Parameter on Cold Solvent Extraction with Water	97
5.3.1	Effect of Extraction Time on Extraction at Room Temperature with Water	98
5.3.2	Effect of Solvent Volume on Extraction at Room Temperature with Water	99
5.3.3	Effect of Particle Size on Extraction at Room Temperature with Water	100
5.4	Validation Method	101
5.4.1	Linearity	102
5.4.2	Limit of Detection and Limit of Quantification	102
5.4.3	Relative Recoveries	103
	5.4.3.1 Aqueous Solvent Extraction with Ultrasonic-Assisted Extraction	103
	5.4.3.2 Aqueous Solvent Extraction with Cold Solvent Extraction	104
5.5	Ultrasonic-Assisted Extraction Application on Cyclodextrin	

	Complexation with Turmeric Rhizomes Oleoresin	105
	5.5.1 Solubility of Curcumin	105
5.6	Characterization	106
	5.6.1 Scanning Electron Microscope	106
	5.6.2 Fourier Transform Infrared Spectroscopy	110
5.7	Summary	114
6	MICROWAVE-ASSISTED EXTRACTION OF CURCUMINOIDS AND CYCLODEXTRIN-COMPLEXATION TURMERIC USING AQUEOUS SOLVENT	116
6.1	Introduction	116
6.2	Effect of Extraction Parameters on Microwave-Assisted Extraction with Water	117
	6.2.1 Effect of Particle Size on Microwave-Assisted Extraction with Water	117
	6.2.2 Effect of Extraction Time on Microwave-Assisted Extraction with Water	119
	6.2.3 Effect of Solvent Volume on Microwave-Assisted Extraction with Water	121
	6.2.4 Effect of Temperature on Microwave-Assisted Extraction with Water	122
	6.2.4.1 Microwave Power	124
6.3	Validation Method	124
	6.3.1 Linearity	125
	6.3.2 Limit of Detection	125
	6.3.3 Relative Recoveries	126
	6.3.3.1 Aqueous Solvent Extraction	126
6.4	Microwave-Assisted Extraction of Cyclodextrin-Complexed Turmeric Oleoresin from Rhizomes of <i>C. domestica</i> Val.	127
	6.4.1 Solubility of Curcumin	127
6.5	Characterization	129
	6.5.1 Scanning Electron Microscope	130

6.5.2	Fourier Transform Infrared Spectroscopy	133
6.6	Summary	137
7	CONCLUSIONS AND RECOMMENDATIONS	138
7.1	Conclusions	138
7.2	Suggestions for Further Study	140
	REFERENCES	142
	Appendice A	158

LIST OF TABLES

TABLE NO.	TITLE	PAGE
2.1	The Properties of α -, β -, and γ -Cyclodextrins (Connors, 1997)	36
3.1	Parameters of ultrasonic-assisted extraction	47
3.2	Parameters of cold solvent extraction	52
3.3	Acceptable value for reproducibility (AOAC)	57
4.1	Polarity of Curcuminoids	64
4.2	Percentage relative recovery and relative standard deviation for turmeric samples with methanol using ultrasonic- assistance extraction	80
4.3	Percentage relative recovery and relative standard deviation for turmeric samples with methanol at room temperature	81
5.1	Validation data of curcuminoids from turmeric sample by ultrasonic-assisted extraction	102
5.2	Percentage relative recovery and relative standard deviation for turmeric samples with ultrasonic-assisted extraction using aqueous solvent	104
5.3	Percentage relative recovery and relative standard deviation for turmeric sample with water at room temperature	104
6.1	Validation data of curcuminoids from turmeric sample using microwave-assisted extraction	125
6.2	Percentage relative recovery and relative standard deviation of turmeric samples with microwave-assisted extraction using aqueous solvent	127

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
2.1	Structures of curcumin and related curcuminoids	13
2.2	Scheme of a conventional Soxhlet extractor (Peter and Naoko, 2013)	16
2.3	UAE device (Katarzyna <i>et al.</i> , 2003)	20
2.4	Basic heat and mass transfer mechanisms in microwave and conventional extraction of natural product (Périno-Issartier <i>et al.</i> , 2011)	23
2.5	Schematic representation of yield versus time in extraction processes. (Raynie, 2000)	24
2.6	A schematic setup for SWE (Rangsriwong <i>et al.</i> , 2008)	29
2.7	Schematic of SFE (Watanabe <i>et al.</i> , 2004)	31
2.8	Schematic of a PLE (Richter <i>et al.</i> , 1996)	32
2.9	Chemical structure of M β -CD	33
2.10	Structures of α -, β - and γ -CDs (Veen <i>et al.</i> , 2000)	34
2.11	Cone shape of CD (Brewster and Loftsson, 2007)	35
2.12	Approximate dimensions of α -, β - and γ -CDs (Szejtli, 2004)	36
2.13	Inclusion complexation between curcumin: M β -CD	38
2.14	Phase solubility diagram (Brewster and Loftsson, 2007)	41
3.1	Schematic representation of UAE setup (Jadhav <i>et al.</i> , 2009)	46
3.2	Schematic representation of MAE setup (http://globalmicrowave) (2013)	48

3.3	Flowchart of experimental procedure for isolation of curcuminoids in <i>C. domestica</i> Val. and inclusion complexation with CD	62
4.1	Relative response factor variations of curcuminoids with ultrasonic amplitude using UAE at room temperature. Extraction conditions: extraction time of 10 min, solvent volume of 10 mL, particle size between 0.06 – 3.35 mm and 2 g of sample with methanol. Chromatographic conditions: mobile phase ACN-H ₂ O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 ×100 mm, 5 µm) and UV detection	65
4.2	Variations of relative response factor of analytes against particle size of the sample using UAE at optimum conditions of extraction at room temperature, solvent volume of 10 mL, ultrasonic amplitude of 100, extraction time of 10 min and 2 g of sample with methanol. Chromatographic conditions: mobile phase ACN-H ₂ O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 ×100 mm, 5 µm) and UV detection	67
4.3	Variations of relative response factor of analytes against various times using UAE at optimum conditions of extraction at room temperature, an ultrasonic amplitude of 100, particle size of 0.30 – 0.60 mm, solvent volume of 10 mL and 2 g of sample with methanol. Chromatographic conditions: mobile phase ACN-H ₂ O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 ×100 mm, 5 µm) and UV detection	68
4.4	Variations of relative response factor of analytes against various solvent volumes with the UAE at optimum conditions of extraction at room temperature, ultrasonic amplitude of 100, particle size of 0.30 - 0.60 mm, extraction time of 10 min and 2 g of sample with methanol. Chromatographic conditions: mobile phase ACN-H ₂ O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 ×100 mm, 5 µm) and UV detection	70
4.5	Variations of relative response factor of analytes against various	

- temperatures using UAE at optimum conditions of extraction of ultrasonic amplitude of 100, particle size of 0.30 - 0.60 mm, extraction time of 10 min and 2 g of sample with methanol. Chromatographic conditions: mobile phase ACN-H₂O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection 72
- 4.6 Variations of relative response factor of analytes against methanol concentration in the extraction solvent using UAE under optimum extraction conditions: extraction temperature of 60°C, ultrasonic amplitude of 100, particle size of 0.30 - 0.60 mm, extraction time of 10 min and 2 g of sample. Chromatographic conditions: mobile phase ACN-H₂O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection 73
- 4.7 Variations of relative response factor of analytes against various extraction times using cold solvent extraction at optimum conditions of extraction at room temperature: solvent volume of 10 mL, particle size of 0.06 - 3.35 mm and 2 g of sample with methanol. Chromatographic conditions: mobile phase ACN-H₂O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection 75
- 4.8 Variations of relative response factor of analytes against various solvent volumes using cold solvent (methanol) extraction. Extraction conditions: extraction at room temperature, solvent volume of 10 mL, particle size of 0.06 - 3.35 mm, extraction time of 60 min and 2 g of sample with methanol. Chromatographic conditions: ACN-H₂O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection 76
- 4.9 Variations of the relative response factor of analytes against various particle sizes using cold solvent extraction. Extraction conditions: extraction at room temperature, solvent volume of 5 mL, extraction time of 60 min and 2 g of sample with methanol.

- Chromatographic conditions: mobile phase ACN-H₂O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection. 78
- 4.10 Chromatogram of a curcumin (C) standard. Chromatographic conditions: mobile phase ACN-H₂O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection 82
- 4.11 Chromatogram of a demethoxycurcumin (DMC) standard. Chromatographic conditions: mobile phase ACN-H₂O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection 82
- 4.12 Chromatogram of a bisdemethoxycurcumin (BDMC) standard. Chromatographic conditions: mobile phase ACN-H₂O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection 83
- 4.13 Chromatogram of a biphenyl standard (Internal standard). Chromatographic conditions: mobile phase ACN-H₂O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection 83
- 4.14 Chromatograms of the mixture of pure standard for bisdemethoxycurcumin (BDMC), demethoxycurcumin (DMC), curcumin (C). Chromatographic conditions: mobile phase ACN-H₂O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection 84
- 4.15 Chromatogram of a biphenyl standard (Internal standard), bisdemethoxycurcumin (BDMC), demethoxycurcumin (DMC), curcumin (C). Chromatographic conditions: mobile phase ACN-H₂O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection 84
- 4.16 Chromatogram of real sample for Biphenyl standard (Internal standard), bisdemethoxycurcumin (BDMC), demethoxycurcumin (DMC), curcumin (C). Chromatographic conditions: mobile phase ACN-H₂O (30:70; v/v), 1 mL/min,

- temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection 85
- 4.17 Combined chromatograms of real sample and standards of curcuminoids. Chromatographic conditions: mobile phase ACN-H₂O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection 85
- 5.1 Variations of relative response factor of analytes against various ultrasonic amplitudes using UAE at room temperature, extraction time of 20 min, solvent volume of 10 mL, particle size between 0.06 - 3.35 mm and 2 g of sample with water. Chromatographic conditions: mobile phase ACN-H₂O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection 90
- 5.2 Variations of relative response factor of analytes against various particle sizes using UAE at room temperature at optimum conditions: solvent volume of 10 mL, ultrasonic amplitude of 100, extraction time of 20 min and 2 g of sample with water. Chromatographic conditions: mobile phase ACN-H₂O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection 92
- 5.3 Variations of relative response factor of analytes against various extraction times using UAE at room temperature at optimum conditions: ultrasonic amplitude of 100, particle size of 0.30 – 0.60 mm, solvent volume of 10 mL and 2 g of sample with water. Chromatographic conditions: mobile phase ACN-H₂O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection 94
- 5.4 Variations of relative response factor of analytes against various solvent volumes with the UAE at room temperature using optimum conditions: ultrasonic amplitude of 100, particle size of 0.30 - 0.60 mm, extraction time of 20 min and 2 g of sample with water. Chromatographic conditions: mobile phase ACN-H₂O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP

	column (4.6 × 100 mm, 5 µm) and UV detection	95
5.5	Variations of relative response factor of analytes against various temperatures using UAE at optimum conditions: ultrasonic amplitude of 100, particle size of 0.30 - 0.60 mm, solvent volume of 10 mL, extraction time of 20 min and 2 g of sample with water. Chromatographic conditions: mobile phase ACN-H ₂ O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection	97
5.6	Variations of relative response factor of analytes against various times using cold solvent extraction at optimum conditions of extraction at room temperature, solvent volume of 10 mL, particle size of 0.06- 3.35 mm and 2 g of sample with water. Chromatographic conditions: mobile phase ACN-H ₂ O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection	98
5.7	Variations of relative response factor of analytes against various solvent volumes using cold solvent extraction at optimum conditions of extraction at room temperature, particle size of 0.06 - 3.35 mm and extraction time of 60 min and 2 g of sample with water. Chromatographic conditions: mobile phase ACN-H ₂ O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection	99
5.8	Variations of relative response factor of analytes against various particle sizes using cold solvent extraction at optimum conditions of extraction at room temperature, solvent volume of 10 mL and extraction time of 60 min and 2 g of sample with water. Chromatographic conditions: mobile phase ACN-H ₂ O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection	101
5.9	Solubility of curcumin as a function of Mβ-CD (a), and plot of linearity graph ranges from 0 - 200 mM Mβ-CD (b), in ethanol-water (20:80; v/v) solution at 30°C, data point is the average of 3 measurements.	105

5.10	SEM photograph (magnification $\times 10000$) of turmeric rhizomes oleoresin (TRO) after extraction using UAE with aqueous solvent	107
5.11	SEM photograph (magnification $\times 5000$) of M β -CD	108
5.12	SEM photograph (magnification $\times 5000$) of turmeric rhizomes oleoresin-methyl- β -cyclodextrin physical mixture method (TRO-M β -CD PM) using UAE with aqueous solvent	108
5.13	SEM photograph (magnification $\times 15000$) of turmeric rhizomes oleoresin-methyl- β -cyclodextrin kneading method (TRO-M β -CD K) using UAE with aqueous solvent	109
5.14	SEM photograph (magnification $\times 25000$) of turmeric rhizomes oleoresin-methyl- β -cyclodextrin co-precipitation method (TRO-M β -CD CP) using UAE with aqueous solvent	109
5.15	FTIR spectrum of turmeric rhizomes oleoresin (TRO) after extraction with UAE using aqueous solvent	111
5.16	FTIR spectrum of turmeric rhizomes oleoresin-methyl- β -cyclodextrin kneading complex (TRO-M β -CD K) using UAE with aqueous solvent	112
5.17	FTIR spectrum of turmeric rhizomes oleoresin-methyl- β -cyclodextrin co-precipitation method (TRO-M β -CD CP) using UAE with aqueous solvent	112
5.18	FTIR spectrum of turmeric rhizomes oleoresin-methyl- β -cyclodextrin physical mixture method (TRO-M β -CD PM) using UAE with aqueous solvent	113
5.19	FTIR spectrum of pure M β -CD	113
6.1	Variations of relative response factor of analytes against particle size using MAE at room temperature at optimum conditions of extraction, solvent volume of 10 mL, extraction time of 4 min and 2 g of sample with water. Chromatographic conditions: mobile phase ACN-H ₂ O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 \times 100 mm, 5 μ m) and UV detection	118
6.2	Variations of relative response factor of analytes with extraction	

	time using MAE at optimum conditions: room temperature, particle size of 0.30 – 0.60 mm, solvent volume of 10 mL and 2 g of sample with water. Chromatographic conditions: mobile phase ACN-H ₂ O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection	120
6.3	Variations of relative response factor of analytes with solvent volume using MAE at optimum conditions; at room temperature, particle size of 0.30 – 0.60 mm, extraction time of 3 min, 2 g of sample and water. Chromatographic conditions: mobile phase ACN-H ₂ O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection	122
6.4	Variations of relative response factor of analytes with temperature using MAE. Extraction conditions: particle size, 0.30 – 0.60 mm; extraction time, 3 min; 2 g of sample and water. Chromatographic conditions: mobile phase ACN-H ₂ O (30:70; v/v), 1 mL/min, temperature 40°C, 254 nm, PFP column (4.6 × 100 mm, 5 µm) and UV detection	123
6.5	Solubility of curcumin as a function of Mβ-CD (a), and plot of linearity graph ranges from 0-200 mM Mβ-CD (b), in ethanol-water (20:80 v/v) solution at 30°C. Each data point is the average of three measurements	128
6.6	SEM image (magnification × 10,000) of turmeric rhizomes oleoresin (TRO) after extraction using MAE with aqueous solvent	130
6.7	SEM image (magnification × 5000) of Mβ-CD	131
6.8	SEM image (magnification × 25,000) of turmeric rhizomes oleoresin-methyl-β-cyclodextrin physical mixture method (TRO-Mβ-CD PM) using MAE with aqueous solvent	131
6.9	SEM image (magnification × 25,000) of turmeric rhizomes oleoresin-methyl-β-cyclodextrin kneading method (TRO-Mβ-CD K) using MAE with aqueous solvent	132
6.10	SEM image (magnification × 25,000) of turmeric rhizomes	

	oleoresin-methyl- β -cyclodextrin co-precipitation method (TRO-M β -CD CP) using MAE with aqueous solvent	132
6.11	FTIR spectrum of turmeric rhizomes oleoresin (TRO) after extraction with MAE using aqueous solvent	134
6.12	FTIR spectrum of turmeric rhizomes oleoresin-methyl- β -cyclodextrin kneading complex (TRO-M β -CD K) using MAE with aqueous solvent	135
6.13	FTIR spectrum of turmeric rhizomes oleoresin-methyl- β -cyclodextrin co-precipitation method (TRO-M β -CD CP) using MAE with aqueous solvent	135
6.14	FTIR spectrum of turmeric rhizomes oleoresin-methyl- β -cyclodextrin physical mixture method (TRO-M β -CD PM) using MAE with aqueous solvent	136
6.15	FTIR spectrum of pure M β -CD	136

LIST OF ABBREVIATIONS

α -CD	-	Alpha-cyclodextrin
ACN	-	Acetonitrile
ASE	-	Accelerated solvent extraction
AOAC	-	Association of analytical communities
ATR	-	Attenuated Total Reflectance
BDMC	-	Bisdemethoxycurcumin
β -CD	-	Beta-cyclodextrin
BP	-	Biphenyl
CD	-	Cyclodextrin
C	-	Curcumin
CP	-	Co-precipitation
DMC	-	Demethoxycurcumin
DMAE	-	Dynamic microwave assisted extraction
FTIR	-	Fourier transform infrared spectroscopy
γ -CD	-	Gamma-cyclodextrin
GC-MS	-	Gas Chromatography-Mass Spectrometry
HS-SPME	-	Head space-solid phase microextraction
HPLC	-	High performance liquid chromatography
HPLC-FD	-	High performance liquid chromatography Flourescence detection
HP β CD	-	Hydroxylpropyl β -cyclodextrin
HP γ CD	-	2-hydroxypropyl- γ -cyclodextrin
IS	-	Internal standard
K	-	Kneading
LOD	-	Limit of detection

LOQ	-	Limit of quantification
M β -CD	-	Methyl- β -cyclodextrin
MAE	-	Microwave-assisted extraction
MAE-HS-SPME	-	Microwave-assisted extraction head space-solid phase microextraction
M β CD-K	-	Methyl Beta cyclodextrin-kneading
M β CD-CP	-	Methyl Beta cyclodextrin-co-precipitation
M β CD-PM	-	Methyl Beta cyclodextrin-physical mixture
PHWE	-	Pressurised hot water extraction
PLE	-	Pressurised liquid extraction
PSE	-	Pressurised solvent extraction
PAHs	-	Polyaromatic hydrocarbons
PCBs	-	Polychlorinated Biphenyls
PFP	-	Penta fluoro phenyl
PM	-	Physical mixture
RSD	-	Relative standard deviation
RP-HPLC	-	Reversed phase high performance liquid chromatography
RP-HPLC-UV	-	Reversed phase high performance liquid chromatography ultraviolet spectroscopy
RR	-	Relative Recovery
RT	-	Retention time
SFC	-	Supercritical fluid chromatography
SFE	-	Supercritical fluid extraction
SWE	-	Subcritical water extraction
SEM	-	Scanning electron microscope
SWE	-	Subcritical or superheated water extraction
SBE β CD	-	Sulfated butyl ethyl β -cyclodextrin sodium salt
SPME	-	Solid phase microextraction
TLC	-	Tin layer chromatography
TRO-M β -CD	-	Turmeric rhizomes oleoresin-methyl β -cyclodextrin
UV	-	Ultraviolet spectroscopy
UAE	-	Ultrasonic-assisted extraction

LIST OF SYMBOLS

α	-	Alpha
β	-	Beta
r^2	-	Coefficients of determination
CO_2	-	Carbon dioxide
$^{\circ}\text{C}$	-	Degree celsius
γ	-	Gamma
g	-	Gram
g mol^{-1}	-	Gram per mole
h	-	Hour
μm	-	Micrometer
μL	-	Micro liter
μgL^{-1}	-	Microgram per liter
min	-	Minutes
mL	-	Milliliter
mm	-	Millimeter
mgL^{-1}	-	Milligram per liter
nm	-	Nanometer
%	-	Percent
pm	-	Picometer
rpm	-	Revolution per minutes
K_c	-	Stability constant
S_o	-	Intrinsic solubility
k	-	Slope

LIST OF APPENDICE

APPENDIX	TITLE	PAGE
A	List of papers published and presented in seminar	158

CHAPTER 1

INTRODUCTION

1.1 Research Background

Turmeric is a member of the Zingiberaceae family native to South Asia, which is a perennial herb with short and thick rhizomes with yellow flesh (Somchit *et al.*, 2002), grown in warm, rainy regions of the world such as India, China, Indonesia, Jamaica and Peru. Its rhizomes are oblong, ovate, perform and often short branched (Jayaprakasha *et al.*, 2002). The plant reaches up to 3 feet (0.9 m) tall and the leaves are dark and medium green in colour with about 1.5 feet (0.45 m) long by 8 inches (20.3 cm) wide (Jayaprakasha *et al.*, 2005). The dry rhizomes of *C. domestica* Val. have been reported to contain 3 - 5% of essential oil and 0.02 - 2.0% aromatic yellow curcuminoids (Began *et al.*, 2000). It exhibits poor to moderate stability in water, oxidized or changes in pH when exposed to light but, has good tinctorial strength. Turmeric oleoresin is the combination of flavours and colour principles obtained from turmeric, however, the principal colouring material in turmeric and its oleoresin are curcumin. Curcumin is orange yellow in colour, crystalline powder, insoluble in water and hexane. It is partially soluble in ether, dichloromethane and freely soluble in ethanol, methanol, acetone and glacial acetic acid (Cheng *et al.*, 1982).

It has been reported that the leguminous plant has a great commercial and medicinal value. *C. domestica* Val. consists of two constituents: phenolic pigment and essential oil. However, the major components of essential oil are ar-turmerone, zingiberene, turmerone and curlone, while that of phenolic pigments are C, DMC and BDMC (Taylor and McDowel, 1992). The yellow pigments are used as colouring agents and the mature rhizomes are ground to give an aromatic yellow powder which is used as spices (Kan, 1997; Anna *et al.*, 2003). Several phenolic compounds are flavonoids and curcuminoids as reported by Jayaprakasha *et al.* (2005). Health benefits of flavonoids are well known and they displayed a remarkable range of biomedical and pharmacological properties that may significantly affect the function of various mammalian cells (Middleton *et al.*, 2000).

Although several active phytochemicals, and high activity profile drugs have been discovered from plants, but the quality and safety related problems of herbal drugs have still been challenged for researchers. The main problems for these drawbacks are the lack of reliable extraction, methodologies and high performance for establishing the purity and standard for the herbal drugs (Huie, 2002). As a result of the above-mentioned factors, herbal medicines have yet to find their way in order to be accepted in the global market. An individual plant may consist of several active phytochemicals existing in abundance along with certain constituents of the low activity profile. There is thus a need in the development of efficient procedures of extraction and analytical techniques with high performance (Smith, 2003) for extracting the phenolic compounds (Beejmohun *et al.*, 2007).

Recent years have shown growing popularity and faith in the use of herbal medicine worldwide. This is attributed to the realisation that modern synthetic drugs have failed to provide curative and guarantee to most of the human diseases with sometime producing side effects and in the end then give more problems than the actual disease. The natural medicine provides a ray of hope through its phytochemicals, which are believed to act in a synergistic manner, providing healing without side effects (Vivekananda *et al.*, 2007).

There has been an increasing interest in newer extraction techniques, in the herbal drug (Nyiredy, 2004). These techniques are amenable to automation and provide shortened extraction times, organic solvent consumption reduction, prevention of pollution in analytical laboratories and reduced sample preparation costs (Poole and Poole, 1996; Wan and Wong, 1996). These conventional methods for the extraction of natural products from plant material for instance Soxhlet, has been widely used not only as a technique for extraction of natural products, but also as a comparison for newer extraction techniques (Luque-Garcia and Luke de Castro, 2004; Sanghi and Kannamkumarath, 2004) which are characterised by the consumption of large volumes of solvent and energy, lengthy extraction procedures, and the potentially deleterious degradation of labile compounds (Kerem *et al.*, 2005).

However, with the problem of consumption of solvent, often in large quantities, toxic and together with time consumption problem. This also resulted in consideration of more new extraction techniques such as microwave-assisted extraction (MAE) (Letellier and Budzinski, 1999; Lopez-Avila, 1999; Camel, 2000), supercritical fluid extraction (SFE) (Hawthorne, 1990; Bowadt and Hawthorne, 1995; Smith, 1999), subcritical water extraction (SWE) or superheated water extraction and accelerated solvent extraction (ASE) (Bjorklund *et al.*, 2000). In addition, ultrasonic-assisted extraction (UAE) as a novel technique for extraction of plant tissue has gained increasing attention (Djilani *et al.*, 2006). A variety of methods for extraction of plant materials have been reported (Ong *et al.*, 2000). The traditional solvent extraction techniques for natural products are mostly based on solvent type and required long extraction time and have low efficiency. Most of these techniques have advantages and disadvantages with regard to solvent volume, extraction time and extraction efficiency as well as working at a high elevated temperature and pressure which improves the extraction process.

Microwave energy has a greater potential for rapid heating of sample and have long appeared in analytical laboratories. Abu-Samra *et al.* (1975) were the first researchers to use a domestic microwave oven in the laboratory in metals analysis

from biological samples. Since then the application of microwave method has been developed for different applications (Sparr and Bjorklund, 2000). But in the last two decades, pharmaceutical and natural products have become interested in using MAE as an alternative to conventional methods of extraction. The use of MAE has created increasing interest in the last couple of years due to its rapid, safe, easy to use and cost effective. The microwave heating allows a better recovery and a better sensitivity than conventional methods and good linearity (Pare, 1991; Pare *et al.*, 1994). In comparison with pressurized solvent extraction or supercritical fluid extraction, MAE can be conducted at atmospheric pressure and therefore can be initiated very rapidly. Furthermore, it offers a significant improvement in terms of analysis time and solvent consumption. Microwaves find their applications not only in plant matrix extraction but also in pesticides, organic pollutants, metals and polymers (Tatke and Jaswal, 2011).

The possibility of the use of ultrasonication process was due to the fact that ultrasonic waves break the cells of the vegetal matrix and the cell's contents are released into the extraction medium (Vinatoru *et al.*, 1997). The enhancement of extraction efficiency of organic compounds by ultrasonic is due to the phenomenon called acoustic cavitations (Toma *et al.*, 2001). When ultrasonic is used in combination with conventional heating, the effect of ultrasonic treatment increased. Ultrasonic also offers a mechanical effect allowing greater penetration of solvents into the sample matrix, increasing the contact surface area between the solid and liquid phase, as a result the solute quickly diffuses from the solid phase to the solvent (Rostagno *et al.*, 2003). Thus, UAE proved to be an inexpensive, simple and efficient alternative to conventional extraction techniques. Ultrasonic has an important role in food engineering due to consumer interest in minimally processed foods and has a wide range of current and future applications in the food industry (Ertugay *et al.*, 2004).

Cyclodextrin (CD) is a well-known material used as the solubilising agent for hydrophobic drugs. The CD is a cyclic oligosaccharides of α -(1,4)-linked D-glucopyranose units arranged in a ring formation containing a hydrophilic outer

surface (Szejtli, 1982). The hydrophobic internal cavity provides the capability to form inclusion complexes with a variety of “guest” hydrophobic larger molecules. Such binding allows CD to increase the water solubility of the oleoresin, which is very hydrophobic and not soluble in water (Zaibunnisa *et al.*, 2009). CDs are thus often used to enhance the solubility of pesticides for agricultural and environmental application (Zhang *et al.*, 2005).

Among all the CDs, the β -CD (β -CD) and its derivatives are most widely used in research and manufacturing due to cost availability and suitable cavity size for most of the common guests (Loftsson and Brewster, 1996a). It has a molecular weight between 200-800 g/mol (Waleczek *et al.*, 2003). Substituent on the CD molecules is likely to influence the interactions between drug and carrier by changing the shape of the CD cavity or altering the charge-charge interactions (Tonnesen *et al.*, 2002).

Several methods have been developed for the determination of curcumin, including using TLC and spectrophotometry (Janssen and Gole, 1984). Reversed-phase HPLC (RP-HPLC) separated on a styrene-divinylbenzene copolymer column using a diode array detector has been reported by Taylor and McDowell (1992). Tonnesen and Karlsen (1983) and Smith and Witoska (1984) reported the use of RP-HPLC separations using ultraviolet spectroscopy and electrochemical detection. The research work was focused on the application of novel extraction methods of MAE and UAE in isolating three major components of phenolic pigments: curcumin, demethoxycurcumin and bisdemethoxycurcumin in *C. domestica* Val.. Methanol and water were used for the optimized with UAE application while water was only used for optimized MAE application. RP-HPLC coupled with UV was used because it offers a more convenient and accurate means of separating and estimating individual curcuminoids. Solubility optimization of aqueous extracts with CD was carried out with MAE and UAE methods.

1.2 Problem Statement

Curcuminoids are natural substances with a lot of biological activities, including antiinflammatory, antioxidant and antitumor. However, their usage in the pharmaceutical field is limited by their aqueous insolubility as well as their isolation with aqueous solvent. These shortcomings have led to the consideration of methods which enhance the aqueous solubility. Despite being well established, conventional techniques for natural products isolation suffer from a few drawbacks due to their high cost, time consuming, less environmentally friendly and less health promoting compatibility. The research explores unconventional MAE and UAE techniques of natural product isolation that are most promising, efficient, cost effective and environmentally friendly and improving the solubility of extracting compound by inclusion complexes with CD.

1.3 Objectives of Study

The research is aimed towards improved extraction of curcuminoids, a group of less water soluble compounds using UAE and MAE methods and the inclusion complexation of CD with curcumin. In line with the major aim the targeted objectives are as follows:

- i. To optimize UAE methods for turmeric compounds using water and methanol.
- ii. To optimize MAE methods for turmeric compounds using water
- iii. To compare the optimized MAE and UAE methods.

- iv. To investigate the application of CD for improving the solubility of curcumin in aqueous extract for both MAE and UAE.
- v. To characterize the inclusion complex of turmeric using Fourier transform infrared spectroscopy (FTIR) and Scanning Electron Microscope (SEM).

1.4 Scope of the Study

The work was centred on the extraction of the major compounds of phenolic pigments from *C. domestica* Val. using MAE and UAE and the analytes were analysed using HPLC with UV detection. Enhancement of the solubility of CD was studied with methyl beta-cyclodextrin (M β -CD) and the inclusion complex was analysed by HPLC-UV and characterized by FTIR and SEM.

1.5 Justification of the Research

Extraction techniques have been widely investigated to obtain such valuable natural compounds from plants for commercialization. However, the quality and safety related problems of natural drugs have still been challenged for researchers. The main reasons for this drawback are the lack of reliable high performance and extraction, and methodologies for establishing the purity and standard for the herbal drugs. As a result of the above mentioned factors, the herbal medicines have still to find their way in order to be accepted in the global market. Thus, there is a need arises for development of newer extraction techniques in the herbal drug which is amenable to automation, with shortened extraction times, reduced organic solvent

consumption, prevention of pollution in analytical laboratories and reducing sample preparation costs. The ultimate goal is total or partial replacement of organic solvent with aqueous one that is cost effective, environmentally friendly and safe. The issues of aqueous turmeric rhizomes oleoresin crude extract solubility are also an area of interest.

1.6 Framework of the Research

The research entails extraction of curcuminoids from turmeric, a member of the Zingiberaceae family. The sample (rhizomes) was chopped into smaller pieces and air dried for two weeks before ground into powder in a mill blender and finally, stored in a sealed plastic container for further analysis. Extraction experiments were carried out using water for both MAE and UAE applications. Methanol was used in UAE application. In UAE, the parameters optimized are irradiation time, particle size, amplitude, solvent volume and temperature for both aqueous and methanol. The irradiation time, solvent volume, particle size and temperature were optimized with MAE application. The crude extract was carefully decanted and rinsed with the solvents. The combined solutions were centrifuged at 3800 rpm for 15 min, and then evaporated to dryness in a water bath heater. The methanol crude was filtered and evaporated using a rotary evaporator. In both cases the concentration in form of percentage yield was estimated using HPLC.

Phase solubility studies were performed using the optimum 20 mg of evaporated turmeric rhizomes oleoresin crude extract. The crude extract were obtained by the optimized condition of UAE and MAE methods was added to a glass bottle each separately containing M β -CD in 5.0 mL of ethanol-water (20:80; v/v), solution at various concentration ranging from 0 – 200 mM. The apparent stability constant, K_C , of curcumin in turmeric oleoresin for the M β -CD inclusion complexes was calculated from the slope and intercepts of the linear segment of the phase

solubility line. The mixing methods in M β -CD-turmeric rhizomes oleoresin inclusion complex used include kneading method, co-precipitation method and physical mixture method. The complexes were characterized using FTIR and SEM.

1.7 Thesis Organization

An overview of related research works on curcumin, extraction methods, conventional and unconventional extraction methods, CD inclusion complexes were presented in Chapter 2. The review provides a commentary on the general significance of the curcumin. It then provides a summary of the key mechanisms involved and aims to provide some background information that can be utilized in subsequent extraction work. The review also summarizes developments of natural product isolation using unconventional methods.

The experimental program is presented in Chapter 3. The apparatus, laboratory equipment, stock solution preparation, HPLC parameter optimization, optimization of extraction parameters, methodology of validation, phase solubility, preparation of inclusion complexation and its characterisation are explained together with technicality in the sample preparation.

Chapter 4 presents UAE with methanol together with cold solvent extraction. The optimized effect of extraction parameters with methanol using both UAE and cold solvent extraction are also presented in this chapter.

Chapter 5 presents the extraction of CD inclusion complexes with turmeric oleoresin using UAE with water. The optimized extraction parameters with aqueous solvent coupled with CD and cold solvent extraction are also described.

Chapter 6 presents MAE with water coupled with CD in order to improve the solubility of curcumin. The optimized effect of extraction parameters with aqueous solvent coupled with CD are also discussed in this chapter.

The overall conclusions and suggestions for further research are presented in Chapter 7.

precise economic evaluation, an additional experiments for establishing large-scale units is necessary.

Although UAE can be used successfully for extraction purposes, it should be borne in mind that ultrasonic conditions, including amplitude used and time can lead to the destruction of bioactive compounds. It is also well known that ultrasonic can lead to the production of free radicals within the cavitation bubbles and in some circumstances, these free radicals can induce undesirable changes and/or destruction of the compounds extracted. This unfavourable reaction can be looked into further in order to guard against destruction of bioactive compounds and formation of free radical due to acoustic cavitations.

This research looked at the solubility of turmeric rhizomes oleoresin as a whole. However, if the main aim is to increase solubility in the individual curcuminoids, there is a need for purification after extraction using any available and suitable method in order to avoid interference of the other compounds before inclusion complexation is carried out. CD as solubilizing agent is an expensive material used for solubility improvement, a further research is needed to understand whether re-usage is possible as a spent material.

REFERENCES

- Abu-Samra, A., Morris, J. S., and Koirtiyohann, S. R. (1975). Wet Ashing of some Biological Samples in a Microwave Oven. *Anal. Chem.* 47 (8), 1475-1477.
- Achat, S., Tomao, V., Madani, K., Chibane, M., Elmaataoui, M., Dangles, O., and Chemat, F. (2012). *Direct Enrichment of Olive oil in Oleuropein by Ultrasound-Assisted Maceration at laboratory and Pilot Scale.*
- Al-Harahshed, M., and Kingman, S. W. (2004). Microwave-Assisted Leaching: A Review. *Hydrometallurgy.* 73, 189–203.
- Anna, C. C. M., Manzan, C. M., Fabio, S., Toniolo, F. S., Bredow, E., and Povh, P. (2003). Extraction of Essential Oil and Pigments from *Curcuma longa* L. by Steam Distillation and Extraction with Volatile Solvents. *J. Agric. Food Chem.* 51, 6802-6807.
- Arunava, G., Topu, B., and Pulok, K. M. (2011). Validated Method for Estimation of Curcumin in Turmeric Powder. *Indian J. Traditional Knowledge.* 10(2), 247-250.
- Ayala, R. S., and Luquede Castro, M. D. (2001). Continuous Subcritical Water Extraction as a Useful Tool for Isolation of Edible Essential Oils. *J. Food Chem.* 75, 109-113.
- Baglole, K. N., Boland, P. G., and Wagner, B. D. (2005). Fluorescence Enhancement of Curcumin upon Inclusion into Parent and Modified Cyclodextrins. *J. Photochem. Photobiol. A. Chem.* 173, 230-237.
- Basile, A., Jimenez-Carmona, M. M., and Clifford, A. A. (1998). Extraction of Rosemary by Superheated Water. *J. Agric. Food Chem.* 46, 5205-5209.
- Beejmohun, V., Fliniaux, O., Grand, E., Lamblin, F., Bensaddek, L., Christen, P., Kovensky, J., Fliniaux, M.-A., and Mesnard, F. (2007). Microwave-Assisted Extraction of the Main Phenolic Compounds in Flaxseed. *Phytochem. Anal.* 18, 275–282.

- Began, G., Goto, M., Kodama, A., and Hirose, T. (2000). Supercritical Carbon dioxide Extraction of Turmeric (*Curcuma longa*). *J. Agric. Chem.* 48, 2189-2192.
- Betz, J. M., Brown, P. N., and Roman, M. C. (2011). Accuracy, Precision and Reliability of Chemical Measurements in Natural products Research. *Fitoterapia.* 82, 44-52.
- Bisht, S., Feldmann, G., Soni, S., Ravi, R., Karikar, K., Maitra, A., and Maitra, A. (2007). Polymeric Nanoparticle-Encapsulated Curcumin ("nanocurcumin"): A Novel Strategy for Human Cancer Therapy. *J. Nanobiotechnol.* 5.
- Bjorklund, E., Bowadt, S., and Nilsson, T. (2000). Pressurised liquid Extraction of Persistent Trends Organic Pollutants in Environmental Analysis. *Trends Anal. Chem.* 19 (7), 434-445.
- Boonkird, S., Phisalaphong, C., and Phisalaphong, M. (2008). Ultrasound-Assisted Extraction of Capsaicinoids from *Capsicum frutescens* on a Lab and Pilot-Plant Scale. *Ultrason. Sonochem.* 15, 1075-1079.
- Bowadt, S., and Hawthorne, S. B. (1995). Supercritical Fluid Extraction in Environmental Analysis. *J. Chromatogr. A.* 703, 549-571.
- Brewstar, M. E., and Loftsson, T. (2007). Cyclodextrins as Pharmaceutical Solubilizers. *Adv. Drugs Del. Rev.* 59, 645-666.
- Brunner, G. (2005). Supercritical Fluids: Technology and Application to Food Processing. *J. Food Eng.* 67, 21-33.
- Cabral, M. H. M. (1994). Structure and Properties of Cyclodextrins Inclusion Complex Formation. *Rev. Port. Farm.* 64, 77-84.
- Camel, V. (2000). Microwave-Assisted Solvent Extraction of Environmental Samples. *Trend Anal. Chem.* 19(4), 229-248.
- Cannavà, C., Cupri, V., Ficarra, P., Guardo, M., Majolino, D., Stancanelli, R., and Venuti, V. (2008). Physicochemical Characterization of Coumestrol/ β -Cyclodextrins Inclusion complexes by UV-Vis and FTIR-ART spectroscopies. *Vib. Spec.* 48, 172-178.
- Cha, K. H., Kang, S. W., Kim, C. Y., Um, B. H., Na, Y. R., and Pan, C. H. (2010). Effect of Pressurized Liquids on Extraction of Antioxidants from *Chlorella vulgaris*. *J. Agric. Food Chem.* 58(8), 4756-4761.
- Chan, C.-H., Yusoff, R., Ngoh, G.-C., and Kung, F. W.-L. (2011). Microwave-Assisted Extractions of Active Ingredients from Plants. *J. Chromatogr. A.* 1218, 6213-6225.

- Chemat, F., Abert-Vian, M., and Zill-e-Huma, Y.-J. (2009). *Microwave Assisted Separations: Green Chemistry in Action*. In: Pearlman J. T., (ed) Green chemistry Research Trends. Nova Science Publishers, New York. 33–62.
- Chemat, S., Lagha, A., Aitamar, H., Bartels, P. V., and Chemat, F. (2004). Comparison of Conventional and Ultrasound-Assisted Extraction of Carvone and Limonene from Caraway Seeds. *Flavour Frag. J.* 19, 188-195.
- Chen, J. J., Tsai, C. S., Hwang, T. L., Shieh, P. C., Chen, J. F., and Sung, P. J. (2010). Sesquiterpenes from the Rhizomes of *Curcuma longa* with Inhibitory Activity on Superoxide Generation and Elastase release by Neutrophils. *J. Food Chem.* 199, 974-980.
- Chen, Y., Xie, M.-Y., and Gong, X.-F. (2007). Microwave-Assisted Extraction used for the Isolation of Total Triterpenoid Saponins from *Ganoderma atrum*. *J. Food Eng.* 81, 162–170.
- Chen, L., Song, D., Tian, Y., Ding, L., Yu, A., and Zhang, H. (2008). Application of on-line Microwave Sample-Preparation Techniques. *Trends Anal. Chem.* 27, 151–159.
- Cheng, K. L., Ueno, K., and Imamura, T. (1982). *Handbook of Organic Analytical Reagents*, CRC press, Boca Raton, Florida, U.S.A. 511.
- Christina, S., Gunter, P. E., Gerald, R., and Jan, F. (2010). A Validated Method for the Quantification of Curcumin in Plasma and Brain tissue by Fast Narrow-Bore High-Performance Liquid Chromatography with Fluorescence Detection. *Anal. Bioanal. Chem.* 397, 1917-1925.
- Chun-Hua, Z., Xian, L., Chong-De, S., and Kun-Song, C. (2011). Ultrasonic Extraction of Flavonoids and Phenolics from Loquat (*Eriobotrya japonica* Lindl.) flowers. *African J. Biotechnol.* 10(25), 5020-5026.
- Chunhui, D., Jie, J., Ning, L., Yingjia, Y., Gengli, D., and Xiangmin, Z. (2006). Fast Determination of Curcumol, Curdione and Germacrone in three species of *Curcuma* Rhizomes by Microwave-Assisted Extraction followed by Headspace Solid Phase Microextraction and Gas Chromatography-Mass Spectrometry. *J. Chromatogr. A.* 1117, 115-120.
- Connors, K. A. (1997). The Stability of Cyclodextrin Complexes in Solution. *Chem. Rev.* 97(5), 1325–1358.
- Davis, M. E., and Brewster, M. E. (2004). Cyclodextrin-Based Pharmaceuticals: Past, Present and Future. *Nat. Rev. Drug Discov.* 3, 1023–1035.

- Dhobi, M., Mandal, V., and Hemalatha, S. (2009). Optimization of Microwave Assisted Extraction of Bioactive Flavonolignan–Silybinin. *J. Chem. Metrl.* 3(1), 13–23.
- Dinh, S. M., DeNuzzio, J. D., and Comfort, A. R. (1999). Intelligent Materials for Controlled Release. *Am. Chem. Soc.* 113-127.
- Djilani, A., Legseir, B., Soulimani, R., Dickob, A., and Younos, C. (2006). New Extraction Technique for Alkaloids. *J. Braz. Chem. Soc.* 17, 518-520.
- Ertugay, M. F., Mustafa, S., and Memnune, S. (2004). Effect of Ultrasound Treatment on Milk Homogenisation and Particle Size Distribution of Fat. *J. Vet. Animal Sci.* 28, 303-308.
- Eskilsson, C. S., and Bjorklund, E. (2000). Analytical-Scale Microwave-Assisted Extraction. *J. Chromatogr. A.* 902, 227-250.
- Eurachem Guide (1998). *The Fitness for Purpose of Analytical Methods: A Laboratory Guide to Method Validation and Related Topics.*
- Faith, P., Mabiki, J., Magadula, R., Mdegela, H., and Resto, D. M. (2013). Optimization of Extraction Conditions and Phytochemical Screening of Root Extract of *Synadenium glaucescens* Pax. *Int. J. Chem.* 5(4), 1916-9698.
- Farre, M., Perez, S., Goncalves, C., Alpendurada, M. F., and Barcelo, D. (2010). Green Analytical Chemistry in the Determination of Organic Pollutants in the Aquatic Environment. *Trends Anal. Chem.* 29(11), 1347-1362.
- Ficarra, R., Tommasini, S., Raneri, D., Calabro, M. L., Di, M. R., Bella, C., Rustichelli, C., Gamberini, M. C., and Ficara, P. (2002). Study of Flavonoids β -Cyclodextrins Inclusion Complexes by NMR, FTIR, DSC, X-ray Investigation. *J. Pharm. Biomed. Anal.* 29, 1005-1014.
- Fromming, K. H., and Szejtli, J. (1994). *Cyclodextrins in Pharmacy.* Kluwer Academic publishers. 46-50, 86, 95-96.
- Garcia-Zubiri, I., Gonzalez-Gaitano, G., Sanchez, M., and Isasi, J. R. (2003). FTIR Study of Dibenzofuran-2-Carboxylic Acid and its Complexes with β -Cyclodextrins. *Vib. Spec.* 33, 205-213.
- Govindarajan, V. S. (1980). Turmeric-Chemistry, Technology and Quality. *CRC Critical Rev. Food Sci. Nutrit.* 12, 199-301.
- Grigonis, D., Venskutonis, P. R., Sivik, B., Sandahl, M., and Eskilsson, C. S. (2005). Comparison of Different Extraction Techniques for Isolation of Antioxidants from Sweet Grass (*Hierchloë odorata*). *J. Supercrit. Fluids.* 33, 223–233.

- Hao, J.-Y., Han, W., Huang, S.-D., Xue, B.-Y., and Deng, X. (2002). Microwave-Assisted Extraction of Artemisinin from *Artemisia annua* L. *Sep. Purif. Technol.* 28(3), 91–196.
- Hartonen, K., Inkala, K., Kangas, M., and Riekkola, M. L. (1997). Extraction of Polychlorinated Biphenyls with Water under Subcritical Conditions. *J. Chromatogr. A.* 785, 219-226.
- Hawthorne, S. B. (1990). Analytical Scale Supercritical Fluid Extraction. *J. Anal. Chem.* 62, 633A- 642A.
- Hawthorne, S. B., Yang, Y., and Miller, D. J. (1994). Extraction of Organic Pollutants from Environmental Solids with Sub and Supercritical Water. *Anal. Chem.* 66, 912-2920.
- Higuchi, T., and Connors, K. A. (1965). Phase Solubility Techniques. *Adv. Anal. Chem. Instrum.* 4, 117-122.
- Himanshu, M., Darshana, M., Mehta, B. K., and Jain, D. C. (2013). Microwave-Assisted Extraction Studies of Target Analyte Artemisinin from Dried Leaves of *Artemisia annua* L. *Organic Chem. Internat.* 1-6.
- Hromadkova, Z., Kovacikova, J., and Ebringerova, A. (1999). Study of the Classical and Ultrasound-Assisted Extraction of the Corn Cob Xylan. *Indian Crops and Prod.* 9, 101-109.
- Hu, Z., Cai, M., and Liang, H. H. (2008). Desirability Function Approach for the Optimization of Microwave-Assisted Extraction of Saikosaponins from *Radix bupleuri*. *Sep. Purif. Technol.* 61(3), 266–275.
- Huie, C. W. (2002). A Review of Modern Sample-Preparation Techniques for the Extraction and Analysis of Medicinal Plants. *Anal. Bioanal. Chem.* 373, 23–30.
- Hui, W., Ligang, C., Yang, X., Qinglei, Z., Xiaopan, Z., Qi, Z., and Lan, D. (2011). Dynamic Microwave-Assisted Extraction Coupled On-line with Clean-up for Determination of Caffeine in Tea. *J. Food Sci. Technol.*
- Ishtiaq, F., Farooq, R., Farooq, A., Siddique, M., Shah, H., Hassan, M. U., and Shaheen, M. A. (2009). Application of ultrasound in pharmaceuticals world. *Appl. Sci. J.* 6, 886-893.
- Jadhav, B.-K., Mahadik, K.-R., and Paradkar, A.-R. (2007). Development and Validation of Improved Reversed Phase-HPLC Method for Simultaneous Determination of Curcumin, Demethoxycurcumin and Bisdemethoxycurcumin. *J. Chromatogr.* 65, 483-488.

- Jadhav, D., Rekha, B. N., Gogate, P. R., and Rathod, V. K. (2009). Extraction of vanillin from vanilla pods: A comparison study of conventional Soxhlet and ultrasound assisted extraction. *J. Food Eng.* 93, 421–426.
- Janssen, A., and Gole, T. (1984). Thin layer Chromatographic Determination of Curcumin from Turmeric. *Chromatographia*. 18, 546-549.
- Jayaprakasha, G. K., Jaganmohan, R. L., and Sakariah, K. K. (2002). An Improved HPLC Method for the Determination of Curcumin, Demethoxycurcumin and Bisdemethoxycurcumin. *J. Agric. Food Chem.* 50, 3668-3672.
- Jayaprakasha, G. K., Jagan, L., Mohan, R., and Sakariah, K. K. (2005). Chemistry and Biological Activities of *Curcuma longa*. *J. Trends Food Sci. Technol.* 16, 533-548.
- Jensen, W. B. (2007). The origin of the Soxhlet Extractor. *J. Chem. Educ.* 84(12), 1913–1914.
- Jian-Bing, J., Xiang-hong, L., Mei-qiang, C., and Zhi-chao, X. (2006). Improvement of Leaching Process of Geniposide with Ultrasound. *Ultrason. Sonochem.* 13, 455-462.
- Joshi, U., Mane, V., and Joshi, S. V. (2009). Comparative Study of Conventional and Microwave Assisted Extraction of Some Indigenous Drugs. *Res. J. Pharm. Tech.* 2(2), 417- 418.
- Kamran, A., Mohd, M., Altaf, A., Mohd, A., Md Nasar, M., and Deepak, S. (2012). Validated HPTLC Analysis Method for Quantification of Variability in Content of Curcumin in *Curcuma longa* L. (Turmeric) Collected from Different Geographical Region of Indian. *Asian pacific J. Tropical Biomed.* S584-S588.
- Kan, W. S. (1997). *Pharmaceutical Botany (9th ed.)* Taipei, Taiwan. National Research Institute of Chinese Medicine. 642-643.
- Katarzyna, W., Sasi, K., Kazimierz, W., and Joseph, A. C. (2003). Environmentally Friendly Sample Treatment for Speciation Analysis by Hyphenated Techniques. *Green Chem.* 5, 250-259.
- Kaufmann, B., and Christen, P. (2002). Recent Extraction Techniques for Natural Products: Microwave-Assisted Extraction and Pressurised Solvent Extraction. *Phytochem. Anal.* 13, 105–113.
- Kerem, Z., German-Shashoua, H., and Yarden, O. (2005). Microwave-Assisted Extraction of Bioactive Saponins from Chickpea (*Cicer arietinum* L). *J. Sci. Food Agric.* 85, 406- 412.

- Khajeh, M., Akbari, M. A. R., and Sanchooli, E. (2009). Application of Doehlert Design in the Optimization of Microwave Assisted Extraction for Determination of Zinc and Copper Incereal Samples using FAAS. *Food Anal. Methods*. 3(3), 133–137.
- Khan, M. K., Albert-Vian, A. S., Fabiano-Tixier, A. S., Dangles, O., and Chemat, F. (2010). Ultrasound Assisted Extraction of Polyphenols (Flavone Glycosides) from Orange (*Citrus sinensis* L.) Peel. *Food Chem.* 119, 851-858.
- Kim, S. M., and Zayas, J. F. (1989). Processing Parameters of Chymosin Extraction by Ultrasound. *J. Food Sci.* 54, 700-703.
- Kovács, Á., Ganzler, K., and Simon-Sarkadi, L. (1998). Microwave-Assisted Extraction of Free Aminoacids from Foods. *Z Lebensm Unters Forsch A*. 207, 26–30.
- Kumudhavalli, M. V., Saravanan, C., Thamizh, M. M., and Jayakar, B. (2011). Analytical Method Development and Validation of Curcumin in Tablet Dosage Form by RP-HPLC Method. *Int. Res. J. Pharm.* 2(1), 233-236.
- Lemos, M. A., Aliyu, M. M., and Hungerford, G. (2012). Observation of the Location and Form of Anthocyanin in Purple Potato using Time-Resolved Fluorescence. *Food Sci. Emerg. Technol.* 16, 61-68.
- Letellier, M., and Budzinski, H. (1999). Microwave-Assisted Extraction of Organic Compounds. *Analisis*. 27, 259-271.
- Li, H., Pordesimo, L., and Weiss, J. (2004). High Intensity Ultrasound-Assisted Extraction of Oil from Soybeans. *Food Res. Int.* 37, 731-738.
- Li, J., Zu, Y.-G., Fu, Y.-J., Yang, Y.-C., Li, S.-M., Li, Z.-N., and Wink, M. (2010). Optimization of Microwave Assisted Extraction of Triterpene Saponins from Defatted Residue of Yellow Horn (*Xanthocerasorbifolia* Bunge.) Kernel and Evaluation of its Antioxidant Activity. *Innov. Food Sci. Emerg. Technol.* 11, 637–664.
- Li-Yao, W., Mian, Z., Chao-Feng, Z., and Zheng-Tao, W. (2007). Diaryl Derivatives from the Root Tuber of *Curcuma longa*. *J. Am. Chem. Soc.* 36, 476- 480.
- Lin, X., Xiao-Kang, L., and Shiro, T. (2010). Curcumin has Bright Prospects for the Treatment of Multiple Sclerosis. *Int. Immunopharmacol.* 11(3), 323-330.
- Loftsson, T. (1995). Effects of Cyclodextrins on the Chemical Stability of Drugs in Aqueous Solutions. *Drug Stability*. 1, 22-23.
- Loftsson, T., and Brewster, M. E. (1996a). Pharmaceutical Applications of Cyclodextrins. 1. Drug Solubilization and Stabilization. *J. Pharm. Sci.* 85, 1017-1025.

- Loftsson, T., Fridriksdóttir, H., and Gudmundsdóttir, K. (1996b). The Effect of Water-Soluble Polymers on Aqueous Solubility of Drugs. *Int. J. Pharmaceut.* 127, 293-296.
- Loftsson, T., and Fridriksdóttir, H. (1998). The Effect of Water-Soluble Polymers on the Aqueous Solubility and Complexing Abilities of β -cyclodextrin. *Int. J. Pharmaceut.* 163, 115-121.
- Loftsson, T., and Másson, M. (2004). The Effects of Water-Soluble Polymers on Cyclodextrins and Cyclodextrin Solubilization of Drugs. *J. Drug Deliv. Sci. Tech.* 14, 35-43.
- Loftsson, T., Hreinsdóttir, H., and Másson, M. (2005a). Evaluation of Cyclodextrin Solubilization of Drugs. *Int. J. Pharmaceut.* 302, 18-28.
- Loftsson, T., Jarho, P., Másson, M., and Jaervinen, T. (2005b). Cyclodextrins in Drug Delivery. *Expert Opin. Drug Deliv.* 2(2), 335-351.
- Loftssona, T., and Duchene, D. (2007). Cyclodextrins and their Pharmaceutical Applications, *Int. J. Pharm.* 329, 1–11.
- Loftsson, T., Hreinsdóttir, H., and Másson, M. (2007). The Complexation Efficiency. *J. Incl. Phenom. Macrocycl. Chem.* 57, 545-552.
- Lopez-Avila, V. (1999). Sample Preparation for Environmental Analysis: Critical Review. *J. Anal. chem.* 29, 195.
- Luque de Castro, M. D., and Garcia-Ayuso, L. E. (1998). Soxhlet Extraction of Solid Materials: an Outdated Technique with a Promising Innovative Future. *Anal. Chim. Acta.* 369, 1–10.
- Luque-Garcia, J. L., and Luque de Castro, M. D. (2004). Focused Microwave Assisted Soxhlet Extraction: Devices and Applications. *Talanta.* 64, 571-577.
- MAE (2013). *Schematic Representation of Setup* (<http://globalmicrowave.org>) ([accessed](#) on 24 October).
- Maing, Y., and Miller, I. (1981). Curcumin-Metal Colour Complexes and Process for Preparing them. *European Patent Application. EP 25, 637 A1.*
- Majors, R. E. (2006). Modern Techniques for the Extraction of Solid Materials: An update. *LC-GC NAM.* 24(7), 648-660.
- Majors, R. E. (2008). Practical Aspects of Solvent Extraction. *LC-GC N. Am.* 26(12), 1158-1166.

- Mandal, V., Mohan. Y., and Hemalath, S. (2007). Microwave Assisted Extraction An Innovative and Promising Extraction Tool for Medicinal Plant Research. *Phcog. Rev.* 1(1), 7–18.
- Mandal, V., Dewanjee, S., Sahu, R., and Mandal, S. C. (2009). Design and Optimization of Ultrasound Assisted Extraction of Curcumin as an Effective Alternative for Conventional Solid- Liquid Extraction of Natural Products. *Nat. Prod. Commun.* 4(1), 95-100.
- Manish, D., Arun, N., and Shahid, H. A. (2012). Comparison of Conventional and Non-Conventional Methods of Extraction of Heartwood of *Pterocarpus marsupium roxb.* *Acta Poloniae Pharm. Drug Res.* 69(3), 475 – 485.
- Martin, E., Verhoef, J. C., and Merkus, F. W. (1998). Efficiency, Safety and Mechanism of Cyclodextrins as Absorption Enhancers in Nasal Delivery of Peptide and Protein Drugs. *J. Drug Target.* 6, 17-36.
- Mason, T. J., Paniwnyk, L., and Lorimer, J. P. (1996). The Uses of Ultrasound in Food Technology. *Ultrason. Sonochem.* 3, S253-S260.
- Middleton, E., Kandaswami, C., and Theoharides, T. C. (2000). The Effects of Plant Flavonoids on Mammalian Cells: Implications for Inflammation, Heart Disease and Cancer. *Pharmacol. Rev.* 52, 673-751.
- Ming, C., Xiangyang, G., Hanhua, L., and Peilong, S. (2013). Microwave-Assisted Extraction and Antioxidant Activity of Staranise Oil from *Illicium verum Hook.* *Int. J. Food Sci. Technol.*
- Muhammad, K. K., Maryline, A.-V., Anne-Sylvie, F.-T., Olivier, D., and Farid, C. (2010). Ultrasound-Assisted Extraction of Polyphenols (Flavanone Glycosides) from Orange (*Citrus sinensis L.*) Peel. *Food Chem.* 119, 851–858.
- Mukne, A. P., and Nagarsenker, M. S. (2004). Traimterene- β -Cyclodextrin Systems Preparation Characterization and in Vivo Evaluation. *J. AAPS Pharm. Sci. Technol.* 5(1), 1-9.
- Mura, P., Zerrouk, N., Faucci, M., Maestrelli, F., and Chemtob, C. (2002). Comparative Study of Ibuprofan Complexation with Amorphous Beta-Cyclodextrin Derivatives in Solution and in the Solid State. *Eur. J. Pharmaceut. Biopharmaceut.* 54, 181-191.
- Nalluri, B. N., Chowdary, K. P. R., Murthy, K. V. R., Hayman, A. R., and Becket, G. (2003). Physiocochemical Characterisation and Dissolution Properties of Nimesulide-Cyclodextrin Binary Systems. *J. AAPS Pharm. Sci. Tech.* 4(1) Article 2.

- Nishiyama, T., Mae, T., Kishida, H., Tsukagawa, M., Mimaki, Y., and Kuroda, M. (2005). Curcuminoids and Sesquiterpenoids in Turmeric (*Curcuma longa* L.) Suppress an Increase in Blood Glucose Level in Type 2 Diabetic KK-A⁺ Mice. *J. Agric. Food Chem.* 53, 959-963.
- Nyiredy, S. (2004). Separation Strategies of Plant Constituents-Current Status. *J. Chromatogr. B.* 812, 35-51.
- Ong, E. S., Woo, S. O., and Yong, Y. L. (2000). Pressurized Liquid Extraction of Berberine and Aristolochic Acids Plants. *J. Chromatogr. A.* 904, 57-64.
- Pare, J. R. J., Belanger, J. M. R., and Stafford, S. S. (1994). Microwave-Assisted Process (MAP), A New Tool for the Analytical Laboratory. *Trends Anal. Chem.* 13, 176-184.
- Pare, J. R. J. (1991). Microwave-Assisted Natural Products Extraction. Patent No. 519, 588 (5,002,784).
- Périno-Issartier, S., Zille-Huma, Y.-J., Abert-Vian, M., and Chemat, F. (2011). Solvent free Microwave-Assisted Extraction of Antioxidants from Sea Buckthorn (*Hippophae rhamnoides*) Foodby-Products. *Food Bioprocess Technol.* 4, 1020–1028.
- Peter, C., and Naoko, S. (2013). Stability of Reference Masses V: UV/Ozone Treatment of Gold and Platinum Surfaces. *Metrologia.* 50, 27–36.
- Pitha, J., Milecki, J., Fales, H., Pannell, L., and Uekama, K. (1986). Hydroxypropyl Beta- Cyclodextrin–Preparation and Characterization–Effects on Solubility of Drugs. *Int. J. Pharmaceut.* 29, 73–82.
- Poole, C. K., and Poole, S. K. (1996). Trends in Extraction of Semi-Volatile Compounds for Environmental Analysis. *Anal. Commun.* 33, 11H-14H.
- Preetha, A., Sherin, G. T., Ajaikumar, B. K., Chitra, S., Kuzhuvelil, B. H., Bokyung, S., Sheeja, T. T., Krishna, M., Indira, K. P., Kallikat, N. R., and Bharat, B. A. (2008). Biological Activities of Curcumin and its Analogues (Gongeners) made by Man and Mother Nature. *Biochem. Pharmacology.* 76, 1590-1611.
- Puvaneswary, S., Sharifah, M., and Yatimah, A. (2010). Synthesis and Characterisation of the Inclusion Complex of Dicationic Ionic Liquid and β -Cyclodextrin. *Int. J. Mol. Sci.* 11, 3675-3685.
- Qureshi, S., Shah, A. H., and Ageel, A. M. (1992). Toxicity Studies on *Alpinia galanga* and *Curcuma longa*. *Planta Medica.* 58, 124–127.

- Rangsiwong, P., Rangkadilok, N., and Shotipruk, A. (2008). Subcritical Water Extraction of Polyphenolic Compounds from *Terminalia chebula* fruits. *J. Sci.* 35(1), 103-108.
- Ratnambal, M. J. (1986). Evaluation of Turmeric Accession for Quality. *Qualities Plantarum*. 36, 243-252.
- Raynie, D. E. (2000). Extraction In: Wilson ID, Adlard ER, Cooke M, Poolie CF (eds) *Encyclopedia of Separation Science*. Academic Press, San Diego.
- Reische, D. W., Lillard, D. A., and Eitenmiller, R. R. (1998). Antioxidants in Food Lipids. In C. C. Ahoh and D. B. Min (Eds.), *Chemistry, Nutrition and Biotechnology*. New York. Marcel Dekker. 423-448.
- Richardson, J. F., and Harker, J. H. (2002). *Particle Technology and Separation Processes* (5th ed.). Oxford: Butterworth-Heinemann.
- Richter, B. E., Jones, B. A., Ezzel, J. L., Porter, N. L., Avdalovic, N., and Pohl, C. (1996). Accelerated Solvent Extraction; A Technique for Sample Preparation. *Anal. Chem.* 68, 1033-1039.
- Rostagno, M. A., Palma, M., and Barroso, C. G. (2003). Ultrasound-Assisted Extraction of Soya Isoflavones. *J. Chromatogr. A*. 1012, 119-128.
- Routray, W., and Orsat, V. (2011). Microwave-Assisted Extraction of Flavonoids: A Review. *Food Bioprocess Technol.* 5(2), 1–16.
- Rouhani, Sh., Alizadeh, N., Salimi, Sh., and Haji-Ghasemi, T. (2009). Ultrasonic Assisted Extraction of Natural Pigments from Rhizomes of *Curcuma longa* L. *Pog. Color Colorants Coat.* 2, 103-113.
- Ruan, G. H., and Li, G. K. J. (2007). The Study on the Chromatographic Fingerprint of *Fructusxanthii* by Microwave Assisted Extraction Coupled with GC-MS. *J. Chromatogr. B.* 850, 241–248.
- Rui, L., Cheng, X., Min, Y., Hui-Fang, L., Xing, Z., and De-An, G. (2011). Qualitative and Quantitative Analysis of Curcuminoids in Herbal Medicines Derived from *Curcuma* Species. *Food Chem.* 126, 1890-1895.
- Saenger, W. (1980). Cyclodextrin-Einschlu-ß-Verbindungen in Forschung and Industrie. *Angew. Chem.* 92, 343-361.
- Sanagi, M. M., Umi, K. A., and Smith, R. M. (1993). Application of Supercritical Fluid Extraction and Chromatography to the Analysis of Turmeric. *J. Chromatogr. Sci.* 31, 20-25.

- Sanghi, R., and Kannamkumarath, S. S. (2004). Comparison of Extraction Methods by Soxhlet, Sonicator and Microwave in the Screening of Pesticide Residue from Solid Matrices. *J. Anal. Chem.* 59, 1032-1036.
- Santos, D. T., Veggi, P. C., and Meireles, M. A. A. (2012). Optimization and Economic Evaluation of Pressurized Liquid Extraction of Phenolic Compounds from Jabuticaba Skins. *J. Food Eng.* 108, 444-452.
- Shah, M. V., and Rohit, M. C. (2013). Novel Techniques for Isolation and Extraction of Phyto-Constituents from Herbal Plants. *Am. J. Phytomed. Clinical Therapeutics.* 3, 338-350.
- Singh, R., Tønnesen, H. H., Vogensen, S. B., Loftsson, T., and Másson, M. (2010). Studies of Curcumin and Curcuminoids. XXXVI. The Stoichiometry and Complexation Constants of Cyclodextrin Complexes as Determined by the Phase-Solubility Method and UV-Vis Titration. *J. Incl. Phenom. Macrocycl. Chem.* 66, 335-348.
- Sivakumar, V., Ravi, V. V., Rao, P. G., and Swaminathan, G. (2007). Studies on the Use of Power Ultrasound in Solid-Liquid Myrobalan Extraction Process. *J. Cleaner Prod.* 15(18), 1815-1820.
- Smith, R. M., and Witoska, B. A. (1984). Comparison of Detectors for the Determination of Curcumin in Turmeric by High Performance Liquid Chromatography. *Analyst.* 109, 259- 261.
- Smith, R. M. (1999). Supercritical Fluids in Separation Science-the Dreams, the Reality and the Future. *J. Chromatogr. A.* 856, 85-115.
- Smith, R. M. (2002). Extraction with Superheated Water: Review. *J. Chromatogr. A.* 975, 31-46.
- Smith, R. M. (2003). Before the Injection-Modern Methods of Sample Preparation for Chromatographic Separation and Analysis. *J. Chromatogr. A.* 1000, 3-27.
- Somchit, M. N., Sulaiman, M. R., Noratunlina, R., and Ahmad, Z. (2002). Hepatoprotective Effects of *Curcuma longa* Rhizomes in Paracetamol-Induced Liver Damage in Rats. In: *A proceedings of the Regional Symposium on Environment and Natural Resources*. Omar, R., Rahman, A., Latif, Z., Lihan, T., and Adam, J. H. (Eds.) 10-11th April KL., Malaysia. 1, 698-702.
- Sparr, E. C., and Bjorklund, E. (2000). Analytical-Scale Microwave-Assisted Extraction. *J. Chromatogr. A.* 902, 227-250.

- Spigno, G., and De Faveri, D. M. (2009). Microwave-Assisted Extraction of Tea Phenols: A phenomenological Study. *J. Food Eng.* 93, 210–217.
- Suslick, K. S., and Price, G. J. (1999). Applications of Ultrasound to Materials Chemistry. *Annu. Rev. Mat. Sci.* 29, 295–326.
- Suslick, K. S., Casadonte, D., Green, M., and Thompson, M. (1987). Effects of High Intensity Ultrasound on Inorganic Solids. *Ultrason.* 25, 56–59.
- Sylvia, D. T., Denise, M. D., and Thomas, A. K. (1995). Extraction of Phenolic Compounds from Fresh Leaves: A Comparison of Methods. *J. Chem. Ecology.* 21(2), 117-125.
- Szente, L., Mikuni, K., Hashimoto, H., and Szejtli, J. (1998). Stabilisation and Solubilisation of Lipophilic Natural Colorants with Cyclodextrins. *J. Incl. Phenom. Mol. Recognit. Chem.* 32, 81-89.
- Szejtli, J. (1982). *Cyclodextrins and their Inclusion Complexes*, Akadeemiai Kiado: Budapest.
- Szejtli, J. (1998). Introduction and General Overview of Cyclodextrin Chemistry. *Chem. Rev.* 98, 1743-1753.
- Szejtli, J. (2004). Past, Present and Future of Cyclodextrin Research. *Pure Appl. Chem.* 76(10), 1825-1845.
- Tatke, P., and Jaiswal, Y. (2011). An Overview of Microwave Assisted Extraction and its Applications in Herbal Drug Research. *J. Med. Plant.* 5(1), 21-31.
- Taverniers, I., De loose, M., and Van Bockstae, E. (2004). Trends in Quality in the Analytical Laboratory. II. Analytical Method Validation and Quality Assurance. *Trends Anal. Chem.* 23, 535-552.
- Taylor, S. J., and McDowell, I. J. (1992). Determination of the Curcuminoid Pigments in Turmeric (*Curcuma domestica* Val.) by Reversed-Phase High Performance Liquid Chromatography. *J. Chromatographia.* 34(1-2), 73-77.
- Teo, C. C., Tan, S. N., Yong, J. W. H., Hew, C. S., and Ong, E. S. (2008). Evaluation of the Extraction Efficiency of Thermally Labile Bioactive Compounds in *Gastrodia Elata* Blume by Pressurized Hot Water Extraction and Microwave Assisted Extraction. *J. Chromatogr. A.* 1182(1), 34-40.
- Their, H. P., and Zeumer, H. (1987). *Manual of Pesticide Residue Analysis*. Weinheim, New York, VCH.
- Thostenson, E. T., and Chou, T. W. (1999). Microwave Processing: Fundamentals and Applications. *Compos. Part A. Appl. Sci.* 30(9), 1055–1071.

- Toma, M., Vinatoru, M., Paniwnyk, L., and Mason, T. J. (2001). Investigation of the Effects of Ultrasound on Vegetal Tissues During Solvent Extraction. *Ultrason. Sonochem.* 8, 137-142.
- Tonnesen, H. H., and Karlsen, J. (1983). High-Performance Liquid Chromatography of Curcumin and Related Compounds. *J. Chromatogr.* 259, 367-371.
- Tonnesen, H. H., and Karlsen, J. (1987). Studies on Curcumin and Curcuminoids. X. The Use of Curcumin as a Formulation Aid to Protect Light-Sensitive Drugs in Soft Gelatine Capsules. *Int. J. Pharm.* 38, 247-249.
- Tønnesen, H. H. (2002). Solubility, Chemical and Photochemical Stability of Curcumin in Surfactant Solutions. *Pharmazie.* 57, 820-824.
- Tonnesen, H. H., Masson, M., and Loftsson, T. (2002). Studies of Curcumin and Curcuminoids. XXVII. Cyclodextrin Complexation: Solubility, Chemical and Photochemical Stability. *Int. J. Pharmaceut.* 244, 127-135.
- Usman, L. A., Hamid, A. A., George, O. C., Ameen, O. M., Muhammad, N. O., Zubair, M. F., and Lawal, A. (2009). Chemical Composition of Rhizome Essential Oil of *Curcuma longa* L. Growing in North Central Nigeria. *World J. Chem.* 4(2), 178-181.
- Vareed, S. K., Kakarala, M., Ruffin, M. T., Crowell, J. A., Normolle, D. P., Djuric, Z., and Brenner, D. E. (2008). Pharmacokinetics of Curcumin Conjugate Metabolites in Healthy Human Subjects. *Cancer Epidemiol. Biomark Prev.* 17, 1411-1417.
- Veen, B. A., Uitdehaag-Josst, C. M., Dijkstra, W. B., and Dijkhuizen, L. (2000). Engineering of Cyclodextrin Glycosyl Transferase Reaction and Product Specificity. *Biochimica and Biophysica Acta.* 1543, 336-360.
- Vijaylata, P., Ajai, P. G., and Bikram, S. (2006). Improved HPTLC Method for Determination of Curcuminoids from *Curcuma longa*. *J. Liq. Chromatogr. Rel. Technol.* 29(6), 877-887.
- Villamiel, M., and De Jong, P. (2000). Inactivation of *Pseudomonas fluorescens* and *Streptococcus thermophilus* in Trypticase SoyBroth and Total Bacteria in Milk by Conventional Heating. *J. Food Eng.* 45, 171-179.
- Vinatoru, M. (2001). An Overview of the Ultrasonically Assisted Extraction of Bioactive Principles from Herbs. *Ultrason. Sonochem.* 8, 303-313.
- Vinatoru, M., Toma, M., Radu, O., Filip, P. I., Lazurca, D., and Mason, T. J. (1997). The Use of Ultrasound for the Extraction of Bioactive Principles from Plant Materials. *Ultrason. Sonochem.* 4, 135-139.

- Vivek, R. Y., Sarasija, S., Kshama, D., and Seema, Y. (2009). Effect of Cyclodextrin Complexation of Curcumin on its Solubility and Antiangiogenic and Anti-inflammatory Activity in Rat Colitis Model. *AAPS Pharm. Sci. Tech.* 10(3), 752-762.
- Vivekananda, M., Yogesh, M., and Hemalatha, S. (2007). Microwave Assisted Extraction – An Innovative and Promising Extraction Tool for Medicinal Plant Research. *Pharmacognosy Rev.* 1(1), 7-18.
- Vivekananda, M., Yogesh, M., and Siva, H. (2008). Microwave Assisted Extraction of Curcumin by Sample–Solvent Dual Heating Mechanism Using Taguchi L9 orthogonal design. *J. Pharm. Biomed. Anal.* 46, 322-327.
- Wakte, P. S., Sachin, B. S., Patil, A. A., Mohato, D. M., Band, T. H., and Shinde, D. B. (2011). Optimisation of Microwave, Ultrasonic and Supercritical Carbon dioxide Assisted Extraction Techniques for Curcumin from *Curcuma longa*. *Sep. Purif. Tech.* 79, 50-55.
- Waleczek, K. J., Marques, C., Hempel, B., and Schmidt, P. C. (2003). Phase Solubility Study of Pure (-)- α -bisabolol and Camomile Essential Oil with β -Cyclodextrin. *Eur. J. Pharmaceut. Biopharmaceut.* 55, 247-251.
- Wan, H. B., and Wong, M. K. (1996). Minimisation of Solvent Consumption in Pesticide Residue Analysis. *J. Chromatogr. A.* 754, 43-47.
- Wang, L. J. (2010). Advances in Extraction of Plant Products in Nutraceutical Processing. In: Pathak Y (ed) *Handbook of Nutraceuticals, II: Scale up, Processing and Automation*. CRC Press/Taylor & Francis, Boca Raton. 15–52.
- Wang, X., Kim, J.-R., Lee, S.-B., Kim, Y.-J., Jung, M. Y., Kwon, H.-W., and Ahn, Y.-J. (2014). Effects of Curcuminoids Identified in Rhizomes of *Curcuma longa* on BACE-1 Inhibitory and Behavioral Activity and Lifespan of Alzheimer's Disease Drosophila Models. *BMC Complementary and Alternative Medicine.* 14, 88.
- Wang, Y.-L., Xi, G.-S., Zheng, Y.-C., and Miao, F.-S. (2010). Microwave-Assisted Extraction of Flavonoids from Chinese Herb *Radix Puerariae* (Ge Gen). *J. Med. Plants Res.* 4 (4), 304-308.
- Wang, Y., You, J., Yu, Y., Qu, C., Zhang, H., and Ding, L. (2008a). Analysis of Ginsenosides in *Panax Ginseng* in High Pressure Microwave-Assisted Extraction. *Food Chem.* 110(1), 161–167.

- Wang, J., Sun, B., Cao, Y., Tian, Y., and Li, X. (2008b). Optimisation of Ultrasound-Assisted Extraction of Phenolic Compounds from Wheat Bran. *Food Chem.* 106, 804-810.
- Wang, L., and Weller, C. L. (2006). Recent Advances in Extraction of Nutraceuticals from Plants. *Trends Food Sci. Technol.* 17, 300–312.
- Wang, Y.-J., Pan, M. H., Cheng, A.-L., Lin, L.-I., Ho, Y.-S., Hsieh, J.-K., and Lin, C.-Y. (1997). Stability of Curcumin in Buffer Solutions and Characterization of its Degradation Products. *J. Pharm. Biomed. Anal.* 15, 1867- 1876.
- Watanabe, T., Tsushima, S., Yamamoto, I., Tomioka, O., Meguro, Y., Nakashima, M., Wada, R., Nagase, Y., and Fukuzato, R. (2004). Recovery of Alkali Salt by Supercritical Fluid Leaching Method using Carbon dioxide Proc. 2nd Int. Symposium on Supercritical Fluid Technology for Energy and Environment Application. 363-366.
- Wu, J., Lin, L., and Chau, F. (2001). Ultrasound-Assisted Extraction of Ginseng Saponins from Ginseng Roots and Cultured Ginseng Cells. *Ultrason. Sonochem.* 8, 347-352.
- Yang, Y., Soren, B., Steven, B. H., and David, J. M. (1995). Subcritical Water Extraction of Polychlorinated Biphenyls from Soil and Sediment. *J. Anal. Chem.* 67, 4571- 4576.
- Yinshi, S., Zhengbo, L., and Jianhua, W. (2011). Ultrasound-Assisted Extraction of Five Sflavores from Iris Tectorum Maxim. *Sep. Purif. Tech.* 78, 49 – 54.
- Zaibunnisa, A. H., Sain, N., Said, M., Illias, R. M., Mustapha, W. A. W., and Hassan, O. (2009). Characterisation of Cyclodextrin Complexes with Turmeric Oleoresin. *J. Food Chem.* 114, 459-465.
- Zaibunnisa, A. H., Siti, R. R., and Nur Ain, A. H. (2011). Stabilisation of Curcumin with γ -Cyclodextrin: Phase Solubility Study and its Characterization. 2nd Int. Conf. Biotechnol. Food Sci. 9-13.
- Zhang, A., Liu, W., Wang, L., and Wen, Y. (2005). Characterization of Inclusion Complexation between Fenoxaprop-p-ethyl and Cyclodextrin. *J. Agric. Food Chem.* 53 (18), 7193-7197.
- Zhang, Z. S., Wang, L. J., Li, D., Jiao, S. S., Chen, X. D., and Mao, Z. H. (2008). Ultrasound-Assisted Extraction of Oil from Flaxseed. *Sep. Purif. Technol.* 62, 192-198.